

Product datasheet for TA812647AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NEDD4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7A1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7A1
Applications: WB

Recommended Dilution: WB 1:500
Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 160-460 of human

NEDD4 (NP_940682) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 140.7 kDa

Gene Name: neural precursor cell expressed, developmentally down-regulated 4, E3 ubiquitin protein

ligase

Database Link: NP 940682

Entrez Gene 4734 Human

P46934



NEDD4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7A1] - TA812647AM

Background: This gene is the founding member of the NEDD4 family of HECT ubiquitin ligases that

function in the ubiquitin proteasome system of protein degradation. The encoded protein contains an N-terminal calcium and phospholipid binding C2 domain followed by multiple tryptophan-rich WW domains and, a C-terminal HECT ubiquitin ligase catalytic domain. It plays critical role in the regulation of a number of membrane receptors, endocytic machinery


components and the tumor suppressor PTEN. [provided by RefSeq, Jul 2016]

Synonyms: NEDD4-1; RPF1

Protein Families: Druggable Genome

Protein Pathways: Endocytosis, Ubiquitin mediated proteolysis

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NEDD4 ([RC212738], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEDD4 (1:500). Positive lysates [LY405051] (100ug) and [LC405051] (20ug) can be purchased separately from OriGene.