

# **Product datasheet for TA812719**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ALDH4A1 Mouse Monoclonal Antibody [Clone ID: OTI1H10]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1H10
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:500

Reactivity: Human, Dog, Rat, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-503 of human

ALDH4A1 (NP\_001154976) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** aldehyde dehydrogenase 4 family member A1

Database Link: NP 001154976

Entrez Gene 212647 MouseEntrez Gene 641316 RatEntrez Gene 612452 DogEntrez Gene 8659

<u>Human</u> <u>P30038</u>



### ALDH4A1 Mouse Monoclonal Antibody [Clone ID: OTI1H10] - TA812719

#### Background:

This protein belongs to the aldehyde dehydrogenase family of proteins. This enzyme is a mitochondrial matrix NAD-dependent dehydrogenase which catalyzes the second step of the proline degradation pathway, converting pyrroline-5-carboxylate to glutamate. Deficiency of this enzyme is associated with type II hyperprolinemia, an autosomal recessive disorder characterized by accumulation of delta-1-pyrroline-5-carboxylate (P5C) and proline. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jun 2009]


Synonyms: ALDH4; P5CD; P5CDh

Protein Families: Druggable Genome


**Protein Pathways:** Alanine, aspartate and glutamate metabolism, Arginine and proline metabolism, Metabolic

pathways

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ALDH4A1 (Cat# [RC204884], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALDH4A1 (Cat# TA812719)(1:500).




Western blot analysis of extracts (35ug) from 10 different cell lines by using anti-ALDH4A1 monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-ALDH4A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA812719) (1:500)



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-ALDH4A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA812719) (1:500)