

Product datasheet for TA812761BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PSMA1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI6C4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6C4
Applications: IF, WB

Recommended Dilution: WB 1:250

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PSMA1 (NP_002777) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 29.4 kDa

Gene Name: proteasome subunit alpha 1

Database Link: NP 002777

Entrez Gene 26440 MouseEntrez Gene 29668 RatEntrez Gene 5682 Human

P25786






Background:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jan 2009]

Synonyms: HC2; HEL-S-275; NU; PROS30
Protein Families: Druggable Genome, Protease


Protein Pathways: Proteasome

Product images:




Western blot analysis of extracts (35ug) from 8 cell lines and 2 tissue lysates by using anti-PSMA1 monoclonal antibody (1:250).






Immunofluorescent staining of Hela cells using anti-PSMA1 antibody ([TA812761]/green, upper left; DAPI/blue, lower left; MERGED, upper right) or Isotype control (MERGED, lower right) (1:100).




Immunofluorescent staining of HEK293T cells using anti-PSMA1 antibody ([TA812761]/green, upper left; DAPI/blue, lower left; MERGED, upper right) or Isotype control (MERGED, lower right) (1:100).



Immunofluorescent staining of NCI-H460 cells using anti-PSMA1 antibody ([TA812761]/green, upper left; DAPI/blue, lower left; MERGED, upper right) or Isotype control (MERGED, lower right) (1:100).





Immunofluorescent staining of PC3 cells using anti-PSMA1 antibody ([TA812761]/green, upper left; DAPI/blue, lower left; MERGED, upper right) or Isotype control (MERGED, lower right) (1:100).