

Product datasheet for TA812775BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Xanthine Oxidase (XDH) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5D8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5D8

Applications: WB

Recommended Dilution: WB 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-280 of human XDH

(NP_000370) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 146.2 kDa

Gene Name: xanthine dehydrogenase

Database Link: NP 000370

Entrez Gene 22436 MouseEntrez Gene 497811 RatEntrez Gene 7498 Human

P47989





Xanthine Oxidase (XDH) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5D8] – TA812775BM

Background: Xanthine dehydrogenase belongs to the group of molybdenum-containing hydroxylases

involved in the oxidative metabolism of purines. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Xanthine dehydrogenase can be converted to xanthine oxidase by reversible sulfhydryl oxidation or by irreversible proteolytic modification. Defects in xanthine dehydrogenase cause xanthinuria, may contribute to adult respiratory stress syndrome, and may potentiate influenza infection through an oxygen metabolite-dependent mechanism. [provided by

RefSeq, Jan 2014]


Synonyms: XAN1; XO; XOR

Protein Families: Druggable Genome

Protein Pathways: Caffeine metabolism, Drug metabolism - other enzymes, Metabolic pathways, Purine

metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY XDH ([RC219402], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XDH (1:500). Positive lysates [LY400135] (100ug) and [LC400135] (20ug) can be purchased separately from OriGene.