

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA812777S

Xanthine Oxidase (XDH) Mouse Monoclonal Antibody [Clone ID: OTI1C10]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI1C10	
Applications:	WB	
Recommended Dilution:	WB 1:500	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG2b	
Clonality:	Monoclonal	
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-280 of human XDH (NP_000370) produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	146.2 kDa	
Gene Name:	xanthine dehydrogenase	
Database Link:	<u>NP_000370</u> <u>Entrez Gene 22436 MouseEntrez Gene 497811 RatEntrez Gene 7498 Human</u> <u>P47989</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Xanthine Oxidase (XDH) Mouse Monoclonal Antibody [Clone ID: OTI1C10] – TA812777S	
Background:	Xanthine dehydrogenase belongs to the group of molybdenum-containing hydroxylases involved in the oxidative metabolism of purines. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Xanthine dehydrogenase can be converted to xanthine oxidase by reversible sulfhydryl oxidation or by irreversible proteolytic modification. Defects in xanthine dehydrogenase cause xanthinuria, may contribute to adult respiratory stress syndrome, and may potentiate influenza infection through an oxygen metabolite-dependent mechanism. [provided by RefSeq, Jan 2014]	
Synonyms:	XAN1; XO; XOR	
Protein Families:	Druggable Genome	
Protein Pathway	: Caffeine metabolism, Drug metabolism - other enzymes, Metabolic pathways, Purine metabolism	

Product images:

170 —	-
130 —	
100 —	
70 —	
55 —	-
40 —	
35 —	
25 —	
15 —	
10 —	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY XDH (Cat# [RC219402], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XDH (Cat# [TA812777])(1:500). Positive lysates [LY400135] (100ug) and [LC400135] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US