

# Product datasheet for TA812826M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Syntaxin 1a (STX1A) Mouse Monoclonal Antibody [Clone ID: OTI2D11]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2D11
Applications: IF, WB

**Reactivity:** WB 1:500, IF 1:100 **Human, Rat, Mouse** 

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human STX1A (NP\_004594) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: syntaxin 1A

Database Link: NP 004594

Entrez Gene 20907 MouseEntrez Gene 116470 RatEntrez Gene 6804 Human

Q16623





Background:


This gene encodes a member of the syntaxin superfamily. Syntaxins are nervous system-specific proteins implicated in the docking of synaptic vesicles with the presynaptic plasma membrane. Syntaxins possess a single C-terminal transmembrane domain, a SNARE [Soluble NSF (N-ethylmaleimide-sensitive fusion protein)-Attachment protein REceptor] domain (known as H3), and an N-terminal regulatory domain (Habc). Syntaxins bind synaptotagmin in a calcium-dependent fashion and interact with voltage dependent calcium and potassium channels via the C-terminal H3 domain. This gene product is a key molecule in ion channel regulation and synaptic exocytosis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009]

Synonyms: HPC-1; P35-1; STX1; SYN1A

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane


**Protein Pathways:** SNARE interactions in vesicular transport

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY STX1A (Cat# [RC209062], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-STX1A (Cat# [TA812826])(1:500).






Western blot analysis of extracts (35ug) from 4 different cell lines by using anti-STX1A monoclonal antibody (1:500).



Western blot analysis of extracts (35ug) from 2 different tissue lysates by using anti-STX1A monoclonal antibody (1:500).



Immunofluorescent staining of SF-268 cells using anti-STX1A antibody ([TA812826]/green, upper left; DAPI/blue, lower left; MERGED, upper right) or Isotype control (MERGED, lower right) (1:100).