

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA812934AM

IL29 (IFNL1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3D4]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI3D4	
Applications:	WB	
Recommended Dilution:	WB 1:500	
Reactivity:	Human	
Host:	Mouse	
lsotype:	lgG2b	
Clonality:	Monoclonal	
Immunogen:	Human recombinant protein fragment corresponding to amino acids 20-200 of human IFNL1 (NP_742152) produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	0.5 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Biotin	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	21.7 kDa	
Gene Name:	interferon, lambda 1	
Database Link:	<u>NP_742152</u> <u>Entrez Gene 282618 Human</u> <u>Q8IU54</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	IL29 (IFNL1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3D4] – TA812934AM	
Background:	This gene encodes a cytokine distantly related to type I interferons and the IL-10 family. This gene, interleukin 28A (IL28A), and interleukin 28B (IL28B) are three closely related cytokine genes that form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by the three genes can be induced by viral infection. Al three cytokines have been shown to interact with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA). [provided by RefSeq, Jul 2008]	
Synonyms:	IL-29; IL29	
Protein Families	Druggable Genome, Secreted Protein, Transmembrane	
Protein Pathway	s: Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway	

Product images:

~ 火

170 —	
130 —	
100 —	
70 —	
55 —	
40 —	
35 —	
25 —	•
15 —	
10 —	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IFNL1 (Cat# [RC219027], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IFNL1 (Cat# [TA812934])(1:500).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US