

## **Product datasheet for TA813032**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **TIGIT Mouse Monoclonal Antibody [Clone ID: OTI9A12]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI9A12

Applications: FC

Recommended Dilution: FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TIGIT (NP\_776160) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 26.1 kDa

**Gene Name:** T-cell immunoreceptor with Ig and ITIM domains

Database Link: NP 776160

Entrez Gene 201633 Human

Q495A1

Background: This gene encodes a member of the PVR (poliovirus receptor) family of immunoglobin

proteins. The product of this gene is expressed on several classes of T cells including follicular B helper T cells (TFH). The protein has been shown to bind PVR with high affinity; this binding is thought to assist interactions between TFH and dendritic cells to regulate T cell dependent

B cell responses. [provided by RefSeq, Sep 2009]







Synonyms: VSIG9; VSTM3; WUCAM


**Protein Families:** Transmembrane

# **Product images:**



Flow cytometric analysis of living 293T cells transfected with TIGIT overexpression plasmid ([RC221447]), Red)/empty vector ([PS100001], Blue) using anti-TIGIT antibody (TA813032). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-TIGIT antibody (TA813032) (1:100).