

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA813050BM

Stefin B (CSTB) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5F2]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5F2
Applications:	IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:500
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-98 of human CSTB (NP_000091) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	11 kDa
Gene Name:	cystatin B
Database Link:	<u>NP_000091</u> <u>Entrez Gene 1476 Human</u> <u>P04080</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Stefin B (CSTB) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5F2] – TA813050BM

Background:The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences.
Some of the members are active cysteine protease inhibitors, while others have lost or
perhaps never acquired this inhibitory activity. There are three inhibitory families in the
superfamily, including the type 1 cystatins (stefins), type 2 cystatins and kininogens. This gene
encodes a stefin that functions as an intracellular thiol protease inhibitor. The protein is able
to form a dimer stabilized by noncovalent forces, inhibiting papain and cathepsins I, h and b.
The protein is thought to play a role in protecting against the proteases leaking from
lysosomes. Evidence indicates that mutations in this gene are responsible for the primary
defects in patients with progressive myoclonic epilepsy (EPM1). One type of mutation
responsible for EPM1 is the expansion in the promoter region of this gene of a
CCCCGCCCGCG repeat from 2-3 copies to 30-78 copies. [provided by RefSeq, Jul 2016]


Synonyms:

CPI-B; CST6; EPM1; EPM1A; PME; STFB; ULD

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CSTB (Cat# [RC203872], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CSTB (Cat# [TA813050])(1:500). Positive lysates [LY424918] (100ug) and [LC424918] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 5 cell lines lysates by using anti-CSTB monoclonal antibody (1:500).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-CSTB mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA813050]) (1:500)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US