

# **Product datasheet for TA813094S**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CD137 (TNFRSF9) Mouse Monoclonal Antibody [Clone ID: OTI1F8]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1F8

Applications: FC

Recommended Dilution: FLOW 1:100

Reactivity: Human

Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TNFRSF9 (NP\_001552) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** tumor necrosis factor receptor superfamily member 9

Database Link: NP 001552

Entrez Gene 3604 Human

Q07011

**Background:** Receptor for TNFSF9/4-1BBL. Possibly active during T cell activation. [UniProtKB/Swiss-Prot


Function]

**Synonyms:** 4-1BB; CD137; CDw137; ILA


Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Cytokine-cytokine receptor interaction




# **Product images:**





Flow cytometric analysis of living 293T cells transfected with TNFRSF9 overexpression plasmid ([RC200664]), Red)/empty vector ([PS100001], Blue) using anti-TNFRSF9 antibody ([TA813094]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-TNFRSF9 antibody ([TA813094]) (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right) using anti-TNFRSF9 antibody ([TA813094]). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).