

Product datasheet for TA813142S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PDZK1 Mouse Monoclonal Antibody [Clone ID: OTI2B4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2B4

Applications: WB

Recommended Dilution: WB 1:250~2000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDZK1 (NP_002605) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 56.9 kDa

Gene Name: PDZ domain containing 1

Database Link: NP 002605

Entrez Gene 5174 Human

O5T2W1




Background:

This gene encodes a PDZ domain-containing scaffolding protein. PDZ domain-containing molecules bind to and mediate the subcellular localization of target proteins. The encoded protein mediates the localization of cell surface proteins and plays a critical role in cholesterol metabolism by regulating the HDL receptor, scavenger receptor class B type 1. Single nucleotide polymorphisms in this gene may be associated with metabolic syndrome, and overexpression of this gene may play a role in drug resistance of multiple myeloma. Pseudogenes of this gene are located on the long arm of chromosome 1. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2011]


Synonyms:

CAP70; CLAMP; NHERF-3; NHERF3; PDZD1

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDZK1 ([RC200444], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDZK1 (1:2000).



Western blot analysis of extracts (35ug) from 1 cell lines lysates by using anti-PDZK1 monoclonal antibody (1:250).