

Product datasheet for TA813205M

CD70 Mouse Monoclonal Antibody [Clone ID: OTI7B2]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI7B2
Applications:	FC
Recommended Dilution:	FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CD70 (NP_001243) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	20.9 kDa
Gene Name:	CD70 molecule
Database Link:	NP_001243 Entrez Gene 970 Human P32970
Background:	Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells. [UniProtKB/Swiss-Prot Function]
Synonyms:	CD27-L; CD27L; CD27LG; LPFS3; TNFSF7; TNLG8A
Protein Families:	ES Cell Differentiation/IPS, Transmembrane



[View online »](#)

Protein Pathways: Cytokine-cytokine receptor interaction


Product images:





Flow cytometric analysis of living 293T cells transfected with CD70 overexpression plasmid ([RC200410]), Red/empty vector ([PS100001], Blue) using anti-CD70 antibody ([TA813205]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).





Flow cytometric analysis of living Raji cells, using anti-CD70 antibody ([TA813205], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).



Flow cytometric analysis of living 786-O cells, using anti-CD70 antibody ([TA813205], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-CD70 antibody ([TA813205]) (1:100).



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h using anti-CD70 antibody ([TA813205]) (Right). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).