

## Product datasheet for TA813212BM

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ICOS Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5G8]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI5G8

Applications: FC

Recommended Dilution: FLOW 1:100~400

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ICOS (NP\_036224) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 20.2 kDa

**Gene Name:** inducible T-cell costimulator

Database Link: NP 036224

Entrez Gene 29851 Human

O9Y6W8

**Background:** The protein encoded by this gene belongs to the CD28 and CTLA-4 cell-surface receptor

family. It forms homodimers and plays an important role in cell-cell signaling, immune

responses, and regulation of cell proliferation. [provided by RefSeq, Jul 2008]

Synonyms: AILIM; CD278; CVID1

**Protein Families:** Secreted Protein, Transmembrane







**Protein Pathways:** 


Cell adhesion molecules (CAMs), Primary immunodeficiency, T cell receptor signaling pathway

## **Product images:**





Flow cytometric analysis of living 293T cells transfected with ICOS overexpression plasmid ([RC207031]), Red)/empty vector ([PS100001], Blue) using anti-ICOS antibody ([TA813212]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:400).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-ICOS antibody ([TA813212]) (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h using anti-ICOS antibody ([TA813212]) (Right). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).