

Product datasheet for TA813221BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD70 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5D8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5D8

Applications: FC

Recommended Dilution: FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD70 (NP_001243) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 20.9 kDa

Gene Name: CD70 molecule

Database Link: NP 001243

Entrez Gene 970 Human

P32970

Background: Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of

costimulated T-cells and enhances the generation of cytolytic T-cells. [UniProtKB/Swiss-Prot

Function]

Synonyms: CD27-L; CD27L; CD27LG; LPFS3; TNFSF7; TNLG8A


Protein Families: ES Cell Differentiation/IPS, Transmembrane






Protein Pathways: Cytokine-cytokine receptor interaction


Product images:




Flow cytometric analysis of living 293T cells transfected with CD70 overexpression plasmid ([RC200410]), Red)/empty vector ([PS100001], Blue) using anti-CD70 antibody ([TA813221]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).




Flow cytometric analysis of living Raji cells, using anti-CD70 antibody ([TA813221], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).





Flow cytometric analysis of living __ cells, using anti-CD70 antibody ([TA813221], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-CD70 antibody ([TA813221]) (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h using anti-CD70 antibody ([TA813221]) (Right). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).