

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA813227M

NFKB1 Mouse Monoclonal Antibody [Clone ID: OTI10D7]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI10D7	
Applications:	WB	
Recommended Dilution:	WB 1:250~500	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG2a	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human NFKB1 (NP_003989) produced in HEK293T cell.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	105.2 kDa	
Gene Name:	nuclear factor kappa B subunit 1	
Database Link:	<u>NP_003989</u> <u>Entrez Gene 18033 MouseEntrez Gene 81736 RatEntrez Gene 4790 Human</u> <u>P19838</u>	




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	ORIGENE NFKB1 Mouse Monoclonal Antibody [Clone ID: OTI10D7] – TA813227M	
Background:	This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Alternative splicing results in multiple transcript variants encoding different isoforms, at least one of which is proteolytically processed. [provided by RefSeq, Feb 2016]	
Synonyms:	CVID12; EBP-1; KBF1; NF-kappa-B1; NF-kappaB; NF-kappabeta; NF-kB; NF-kB1; NFkappaB; NFKB-p50; NFKB-p105	
Protein Families:	Druggable Genome, Transcription Factors	
Protein Pathways: Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Cytosolic DNA-ser pathway, Epithelial cell signaling in Helicobacter pylori infection, MAPK signaling path Metabolic pathways, Neurotrophin signaling pathway, NOD-like receptor signaling pa Pancreatic cancer, Pathways in cancer, Prostate cancer, RIG-I-like receptor signaling pa Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pa		

Product images:

170 —	
130 —	-
100 —	
70 —	-
55 —	
40 —	
35 —	
25 —	
15 —	
10 —	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NFKB1 ([RC208384], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NFKB1 (1:500).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Western blot analysis of extracts (35ug) from Hela cells by using anti-NFKB1 monoclonal antibody (1:250).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US