

## Product datasheet for TA813265BM

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD47 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2A10]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI2A10

**Applications:** FC

Recommended Dilution: FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD47 (NP\_001768) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 35.21 kDa

Gene Name: CD47 molecule

Database Link: NP 001768

Entrez Gene 961 Human

Q08722

**Background:** This gene encodes a membrane protein, which is involved in the increase in intracellular

calcium concentration that occurs upon cell adhesion to extracellular matrix. The encoded protein is also a receptor for the C-terminal cell binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This gene has broad tissue distribution, and is reduced in expression on Rh erythrocytes. Alternatively spliced transcript

variants have been found for this gene. [provided by RefSeq, Jul 2010]






Synonyms: IAP; MER6; OA3


**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** ECM-receptor interaction


## **Product images:**




Flow cytometric analysis of living 293T cells transfected with CD47 overexpression plasmid ([RC218813]), Red) using anti-CD47 antibody ([TA813265]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).




Flow cytometric analysis of living Jurkat cells, using anti-CD47 antibody ([TA813265], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).




Flow cytometric analysis of living Raji cells, using anti-CD47 antibody ([TA813265], Red), compared to an isotype control (green) (1:100).



Flow cytometric analysis of living Ramos cells, using anti-CD47 antibody ([TA813265], Red), compared to an isotype control (green) (1:100).



Flow cytometric analysis of living K562 cells, using anti-CD47 antibody ([TA813265], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).



Flow cytometric analysis of living PBMCs, using anti-CD47 antibody ([TA813265], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).