

# **Product datasheet for TA813313S**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### YAP1 Mouse Monoclonal Antibody [Clone ID: OTI9C2]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI9C2

Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 347-504 of human YAP1

(NP\_001123617) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 54.3 kDa

Database Link: NP 001123617

Entrez Gene 22601 Mouse

**Background:** This gene encodes a downstream nuclear effector of the Hippo signaling pathway which is

involved in development, growth, repair, and homeostasis. This gene is known to play a role in the development and progression of multiple cancers as a transcriptional regulator of this signaling pathway and may function as a potential target for cancer treatment. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by

RefSeq, Aug 2013]


Synonyms: COB1; YAP; YAP2; YAP65; YKI






**Protein Families:** Druggable Genome

# **Product images:**





Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and YAP1-Knockout HeLa cells (KO, Cat# [LC832485]) were separated by SDS-PAGE and immunoblotted with anti-YAP1 monoclonal antibody [TA813313] (1:1000). Then the blotted membrane was stripped and reprobed with anti-PCNA antibody as a loading control.

Western blot analysis of extracts (35ug) from 5 cell lines lysates by using anti-YAP1 monoclonal antibody (1:1000).