

# **Product datasheet for TA813404S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HSP27 (HSPB1) Mouse Monoclonal Antibody [Clone ID: OTI8C7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI8C7

Applications: WB

Recommended Dilution: WB 1:500-1000

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human HSPB1 (NP\_001531) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 22.8 kDa

**Gene Name:** heat shock protein family B (small) member 1

Database Link: NP 001531

Entrez Gene 3315 Human

P04792

**Background:** The protein encoded by this gene is induced by environmental stress and developmental

changes. The encoded protein is involved in stress resistance and actin organization and translocates from the cytoplasm to the nucleus upon stress induction. Defects in this gene are a cause of Charcot-Marie-Tooth disease type 2F (CMT2F) and distal hereditary motor

neuropathy (dHMN). [provided by RefSeq, Oct 2008].







**Synonyms:** CMT2F; HEL-S-102; HMN2B; HS.76067; Hsp25; HSP27; HSP28; SRP27

**Protein Pathways:** MAPK signaling pathway, VEGF signaling pathway

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HSPB1 ([RC201800], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSPB1.(1:1000)



Western blot analysis of extracts (35ug) from 3 cell lines lysates by using anti-HSPB1 monoclonal antibody. (1:500)