

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA813473BM

IL2 Receptor alpha (IL2RA) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2A5]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI2A5
Applications:	FC
Recommended Dilution:	Flow 1:100
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human IL2RA (NP_000408) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	30.82 kDa
Gene Name:	interleukin 2 receptor subunit alpha
Database Link:	<u>NP_000408</u> <u>Entrez Gene 3559 Human</u> <u>P01589</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	IL2 Receptor alpha (IL2RA) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2A5] – TA813473BM
Background:	The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolyisis. Alternately-spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. Mutations in this gene are associated with interleukin 2 receptor alpha deficiency.[provided by RefSeq, Nov 2009]
Synonyms:	CD25; IDDM10; IL2R; IMD41; p55; TCGFR
Protein Families	: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane
Protein Pathway	vs: Cytokine-cytokine receptor interaction, Endocytosis, Hematopoietic cell lineage, Jak-STAT signaling pathway

Product images:





Flow cytometric analysis of living 293T cells transfected with IL2RA overexpression plasmid ([RC215768]), Red)/empty vector ([PS100001], Blue) using anti-IL2RA antibody ([TA813473]). Cells incubated with a non-specific antibody (Green) were used as isotype control. <code>D1D100</code>



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-IL2RA antibody ([TA813473]).(1:100).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h using anti-IL2RA antibody ([TA813473])(Right). Cells incubated with a non-specific antibody (Left) were used as isotype control.(1:100).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US