

## **Product datasheet for TA813497S**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## MICB Mouse Monoclonal Antibody [Clone ID: OTI1C4]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI1C4

Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human Host: Mouse

Isotype: lgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MICB (NP\_005922) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 42.4 kDa

**Gene Name:** MHC class I polypeptide-related sequence B

Database Link: NP 005922

Entrez Gene 4277 Human

Q29980





Background: This gene encodes a heavily glycosylated protein which is a ligand for the NKG2D type II

receptor. Binding of the ligand activates the cytolytic response of natural killer (NK) cells, CD8 alphabeta T cells, and gammadelta T cells which express the receptor. This protein is stress-induced and is similar to MHC class I molecules; however, it does not associate with beta-2-microglobulin or bind peptides. Alternative splicing results in multiple transcript variants.


[provided by RefSeq, Jan 2014]

Synonyms: PERB11.2

**Protein Families:** Druggable Genome

Protein Pathways: Natural killer cell mediated cytotoxicity

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MICB/MICA ([RC222315]/[RC204447], Middle/Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MICB.(1:1000)