

Product datasheet for TA813499

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MICB Mouse Monoclonal Antibody [Clone ID: OTI3E6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3E6

Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MICB (NP_005922) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42.4 kDa

Gene Name: MHC class I polypeptide-related sequence B

Database Link: NP 005922

Entrez Gene 4277 Human

Q29980



Background:


This gene encodes a heavily glycosylated protein which is a ligand for the NKG2D type II receptor. Binding of the ligand activates the cytolytic response of natural killer (NK) cells, CD8 alphabeta T cells, and gammadelta T cells which express the receptor. This protein is stress-induced and is similar to MHC class I molecules; however, it does not associate with beta-2-microglobulin or bind peptides. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014]

Synonyms: PERB11.2

Protein Families: Druggable Genome

Protein Pathways: Natural killer cell mediated cytotoxicity

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MICB/MICA ([RC222315]/[RC204447], Middle/Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MICB.(1:1000)