

Product datasheet for TA813512M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

iNOS (NOS2) Mouse Monoclonal Antibody [Clone ID: OTI1A1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A1
Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 539-970 of human

Nos2 (NP_000616) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 130.9 kDa

Gene Name: nitric oxide synthase 2

Database Link: NP 000616

Entrez Gene 4843 Human

P35228

Background: Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes,

including neurotransmission and antimicrobial and antitumoral activities. This gene encodes a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. Three related pseudogenes are located within the

Smith-Magenis syndrome region on chromosome 17. [provided by RefSeq, Jul 2008].



Synonyms: HEP-NOS; INOS; NOS; NOS2A


Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Arginine and proline metabolism,

Calcium signaling pathway, Long-term depression, Metabolic pathways, Pathways in cancer,

Small cell lung cancer

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NOS2 ([RC211819], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NOS2.(1:1000)