

Product datasheet for TA813551M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MAP2 Mouse Monoclonal Antibody [Clone ID: OTI5F8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5F8

Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1507-1827 of human

MAP2 (NP 002365) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 199.3 kDa

Gene Name: microtubule associated protein 2

Database Link: NP 002365

Entrez Gene 17756 MouseEntrez Gene 4133 Human

P11137

MAP2 Mouse Monoclonal Antibody [Clone ID: OTI5F8] - TA813551M


Background: This gene encodes a protein that belongs to the microtubule-associated protein family. The

proteins of this family are thought to be involved in microtubule assembly, which is an essential step in neurogenesis. The products of similar genes in rat and mouse are neuron-specific cytoskeletal proteins that are enriched in dentrites, implicating a role in determining and stabilizing dentritic shape during neuron development. A number of alternatively spliced variants encoding distinct isoforms have been described. [provided by RefSeq, Jan 2010].

Synonyms: MAP-2; MAP2A; MAP2B; MAP2C

Protein Families: Adult stem cells, Druggable Genome, ES Cell Differentiation/IPS

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAP2 ([RC216775], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAP2.(1:1000)