

## Product datasheet for TA813675BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CD28 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4G8]

### **Product data:**

**Product Type: Primary Antibodies** 

Clone Name: OTI4G8

**Applications:** 

Recommended Dilution: WB 1:500

Reactivity: Human

Host: Mouse Isotype: lgG1

Clonality: Monoclonal

Full length human recombinant protein of human CD28 (NP\_006130) produced in HEK293T Immunogen:

PBS (pH 7.3) containing 1% BSA, 50% glycerol. Formulation:

Concentration: 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

HRP Conjugation:

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt.

**Predicted Protein Size:** 25.1 kDa

Gene Name: CD28 molecule

Database Link: NP 006130

Entrez Gene 940 Human

P10747

Background: The protein encoded by this gene is essential for T-cell proliferation and survival, cytokine

> production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]

Synonyms: Tp44






**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Allograft rejection, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Graft-

versus-host disease, Systemic lupus erythematosus, T cell receptor signaling pathway, Type I

diabetes mellitus, Viral myocarditis

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD28 ([RC211318], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD28.(1:500)