

Product datasheet for TA813801

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NSE (ENO2) Mouse Monoclonal Antibody [Clone ID: OTI3C1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C1
Applications: IHC

Recommended Dilution: IHC 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 188-293 of human NSE

(NP_001966) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 47.3 kDa

Gene Name: enolase 2

Database Link: NP 001966

Entrez Gene 13807 MouseEntrez Gene 24334 RatEntrez Gene 2026 Human

P09104

Background: This gene encodes one of the three enclase isoenzymes found in mammals. This isoenzyme,

a homodimer, is found in mature neurons and cells of neuronal origin. A switch from alpha enolase to gamma enolase occurs in neural tissue during development in rats and primates.

[provided by RefSeq, Jul 2008]







Synonyms: HEL-S-279; NSE

Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways, RNA degradation

Product images:



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-NSE mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 3 min, TA813801)



Immunohistochemical staining of paraffinembedded Human adult brain tissue within the normal limits using anti-NSE mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120 oC for 3 min, TA813801)