

## **Product datasheet for TA814175M**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## P Glycoprotein (ABCB1) Mouse Monoclonal Antibody [Clone ID: OTI2A2]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI2A2

Applications: IHC

Recommended Dilution: IHC 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MDR-1/ABCB1 (NP\_000918) produced in

E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 141.3 kDa

**Gene Name:** ATP binding cassette subfamily B member 1

Database Link: NP 000918

Entrez Gene 170913 RatEntrez Gene 5243 Human

P08183





#### Background:


The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance. The protein encoded by this gene is an ATP-dependent drug efflux pump for xenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier. Mutations in this gene are associated with colchicine resistance and Inflammatory bowel disease 13. Alternative splicing and the use of alternative promoters results in multiple transcript variants. [provided by RefSeq, Feb 2017]

**Synonyms:** ABC20; CD243; CLCS; GP170; MDR1; p-170; P-GP; PGY1

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane


**Protein Pathways:** ABC transporters

# **Product images:**



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-MDR-1/ABCB1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-MDR-1/ABCB1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.