

## Product datasheet for **TA814611**

### Glypican 3 (GPC3) Mouse Monoclonal Antibody [Clone ID: OTI1G5]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1G5
Applications:	IHC
Recommended Dilution:	IHC 1:3500
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment of human GPC3 (NP_001158089) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1.00mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	68.41 kDa
Gene Name:	glypican 3
Database Link:	<a href="#">NP_001158089</a> <a href="#">Entrez Gene 2719 Human</a> <a href="#">P51654</a>



[View online »](#)

**Background:**


Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26, and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi-Behmel syndrome, also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2009]

**Synonyms:**


DGSX; GTR2-2; MXR7; OCI-5; SDYS; SGB; SGBS; SGBS1

**Protein Families:**


Druggable Genome

**Product images:**

IHC staining of FFPE human liver tissue with in the normal limits using anti-GPC3 mouse monoclonal antibody (TA814611) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



IHC staining of FFPE human liver cancer tissue using anti-GPC3 mouse monoclonal antibody (TA814611) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



IHC staining of FFPE human placenta tissue with in the normal limits using anti-GPC3 mouse monoclonal antibody (TA814611) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.