

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA815571

KIR3DL1 Mouse Monoclonal Antibody [Clone ID: OTI1A7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1A7
Applications:	WB
Recommended Dilution:	WB 1:500
Reactivity:	Human
Host:	Mouse
lsotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment of Human KIR3DL1 (NP_037421) produced in Ecoli.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.
Stability:	1 Year
Predicted Protein Size:	49.1kDa
Gene Name:	killer cell immunoglobulin like receptor, three lg domains and long cytoplasmic tail 1
Database Link:	<u>Entrez Gene 3811 Human</u> <u>P43629</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

STATES CONTRACT CONTRACTOR CONTRA

Background:Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed
by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly
homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb
leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among
haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3,
KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular
immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S)
cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory
signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR
proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the
TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for
several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to
play an important role in regulation of the immune response. [provided by RefSeq, Jul 2008]

Product images:



Figure A, Western blot analysis of overexpressed lysates (25ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC) and 13 different KIR proteins using anti-KIR3DL1 antibody TA815571 (1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000).From the results,TA815571 only recognize KIR3DL1 protein.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US