

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA815646

ALKBH5 Mouse Monoclonal Antibody [Clone ID: OTI5B4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5B4
Applications:	WB
Recommended Dilution:	WB 1:5000
Reactivity:	Human, Mouse
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment of Human ALKBH5 (NP_060228) produced in Ecoli.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	51.4KD
Gene Name:	alkB homolog 5, RNA demethylase
Database Link:	<u>NP 060228</u>
	<u>Entrez Gene 54890 Human</u> <u>Q6P6C2</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Scrigene ALKBH5 Mouse Monoclonal Antibody [Clone ID: OTI5B4] – TA815646

Background:Dioxygenase that demethylates RNA by oxidative demethylation: specifically demethylates
N(6)-methyladenosine (m6A) RNA, the most prevalent internal modification of messenger
RNA (mRNA) in higher eukaryotes (PubMed:23177736, PubMed:24489119, PubMed:24616105,
PubMed:24778178). Can also demethylate N(6)-methyladenosine in single-stranded DNA (in
vitro) (PubMed:24616105). Requires molecular oxygen, alpha-ketoglutarate and iron
(PubMed:21264265, PubMed:23177736, PubMed:24489119, PubMed:24616105,
PubMed:24778178). Demethylation of m6A mRNA affects mRNA processing and export
(PubMed:23177736). Required for the late meiotic and haploid phases of spermatogenesis by
mediating m6A demethylation in spermatocytes and round spermatids: m6A demethylation
of target transcripts is required for correct splicing and the production of longer 3' UTR
mRNAs in male germ cells (By similarity).[UniProtKB/Swiss-Prot Function]

Synonyms: ABH5; OFOXD; OFOXD1

Product images:




Figure A, Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC) , human ALKBH5 plasmid ([RC209110], hALKBH5), mouse ALKBH5 plasmid ([MR206184], mALKBH5) using anti-ALKBH5 antibody TA815646(1:5000). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)

Western blot analysis of extracts (50ug per lane) from 6 cell lines lysates by using anti-ALKBH5 monoclonal antibody(TA815646, 1:1000)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US