

Product datasheet for TA815852

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Inhibin beta E chain (INHBE) Mouse Monoclonal Antibody [Clone ID: OTI14B5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI14B5

Applications: WB

Reactivity: WB 1:500
Host: WB 0:500
WB 1:500
WB 1:500

Isotype: lgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment of Human INHBE (NP_113667) produced in Ecoli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: 1 Year

Predicted Protein Size: 38.6kDa

Gene Name: inhibin beta E subunit

Database Link: Entrez Gene 83729 Human

P58166

Background: This gene encodes a member of the TGF-beta (transforming growth factor-beta) superfamily

of proteins. The encoded preproprotein is proteolytically processed to generate an inhibin beta subunit. Inhibins have been implicated in regulating numerous cellular processes including cell proliferation, apoptosis, immune response and hormone secretion. This gene may be upregulated under conditions of endoplasmic reticulum stress, and this protein may inhibit cellular proliferation and growth in pancreas and liver. [provided by RefSeq, Sep 2016]





Product images:





Figure A, Western blot analysis of overexpressed lysates (25ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], lane 1), human INHBE plasmid ([RC201426], lane 2), mouse INHBE plasmid ([MR205316], lane 3) using anti-INHBE antibody TA815852 (1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000).



Western blot analysis of extracts (50ug per lane) from HepG2 lysates by using anti-INHBE monoclonal antibody (TA815852, 1:500).