

## Product datasheet for **TA890002M**

### NF-kB p65 (RELA) Rabbit Polyclonal Antibody

#### Product data:


Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide conjugated to KLH derived from within residues 100 - 170 of Human RELA.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from the immunized serum by affinity chromatography (Protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	60 kDa
Gene Name:	RELA proto-oncogene, NF-kB subunit
Database Link:	<a href="#">NP_068810</a> <a href="#">Entrez Gene 19697 Mouse</a> <a href="#">Entrez Gene 309165 Rat</a> <a href="#">Entrez Gene 5970 Human</a> <a href="#">Q04206</a>
Background:	NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011]
Synonyms:	NFKB3; p65


[View online »](#)


**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor signaling pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Cytosolic DNA-sensing pathway, Epithelial cell signaling in Helicobacter pylori infection, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, RIG-I-like receptor signaling pathway, Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway


### Product images:




Western blot analysis of HEK293 cell lysate (35ug) by using Rabbit polyclonal anti-RELA antibody at 1:2000 dilution.




Western blot analysis of HeLa cell lysate (35ug) by using Rabbit polyclonal anti-RELA antibody at 1:2000 dilution.




Western blot analysis of HT29 cell lysate (35ug) by using Rabbit polyclonal anti-RELA antibody at 1:2000 dilution




Western blot analysis of MCF7 cell lysate (35ug) by using Rabbit polyclonal anti-RELA antibody at 1:2000 dilution.




Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and RELA-Knockout HeLa cells (KO, Cat# [LC810201]) were separated by SDS-PAGE and immunoblotted with anti-RELA monoclonal antibody [TA890002], (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90AB1 antibody ([TA500494]) as a loading control.




Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using Rabbit polyclonal anti-RELA antibody at. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human adult heart tissue using Rabbit polyclonal anti-RELA antibody at. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human melanoma tissue using Rabbit polyclonal anti-RELA antibody at. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human testicular cancer tissue using Rabbit polyclonal anti-RELA antibody at. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.