

Product datasheet for TA890030M

PKC delta (PRKCD) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB: 1:500~1:1000 Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human PRKCD

Formulation: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Concentration: 2.18mg/ml

Purification: Purified from the immunized serum by affinity chromatography (Protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 77.3 kDa

Gene Name: protein kinase C delta

Database Link: NP 997704

Entrez Gene 18753 MouseEntrez Gene 170538 RatEntrez Gene 5580 Human

Q05655



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play distinct roles in cells. The protein encoded by this gene is one of the PKC family members. Studies both in human and mice demonstrate that this kinase is involved in B cell signaling and in the regulation of growth, apoptosis, and differentiation of a variety of cell types. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq, Jul 2008]

Synonyms: ALPS3; CVID9; MAY1; nPKC-delta; PKCD


Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Chemokine signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated

phagocytosis, GnRH signaling pathway, Neurotrophin signaling pathway, Tight junction, Type


II diabetes mellitus, Vascular smooth muscle contraction

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PRKCD (Cat# [RC208127], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PRKCD antibody (Cat# [TA890030]). Positive lysates [LY403724] (100ug) and [LC403724] (20ug) can be purchased separately from OriGene.





Western blot analysis of extracts (35ug) from different cell lines or tissues by using anti-PRKCD rabbit polyclonal antibody ([TA890030]).