

## **Product datasheet for TA890126M**

## **HIRA Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500~2000

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

**Isotype:** lgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human HIRA

**Formulation:** PBS with 0.02% sodium azide, 50% glycerol, pH7.3

**Concentration:** 1.65 mg/ml

**Purification:** Purified from the immunized serum by affinity chromatography (Protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 111.7 kDa

**Gene Name:** histone cell cycle regulator

Database Link: NP 003316

Entrez Gene 15260 MouseEntrez Gene 363849 RatEntrez Gene 7290 Human

P54198

**Background:** This gene encodes a histone chaperone that preferentially places the variant histone H3.3 in

nucleosomes. Orthologs of this gene in yeast, flies, and plants are necessary for the

formation of transcriptionally silent heterochomatin. This gene plays an important role in the formation of the senescence-associated heterochromatin foci. These foci likely mediate the irreversible cell cycle changes that occur in senescent cells. It is considered the primary candidate gene in some haploinsufficiency syndromes such as DiGeorge syndrome, and insufficient production of the gene may disrupt normal embryonic development. [provided

by RefSeq, Jul 2008]

Synonyms: DGCR1; TUP1; TUPLE1



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




**Protein Families:** 

**Transcription Factors** 

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HIRA (Cat# [RC208049], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HIRA rabbit polyclonal antibody (Cat# [TA890126]).



Western blot analysis of extracts (35ug) from different cell lines and tissues by using anti-HIRA rabbit polyclonal antibody.