

Product datasheet for TA890156M

XPC Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500~2000

Reactivity: Human, Rat, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human XPC

Formulation: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Concentration: 2.55 mg/ml

Purification: Purified from the immunized serum by affinity chromatography (Protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 105.8 kDa

Gene Name: XPC complex subunit, DNA damage recognition and repair factor

Database Link: NP 004619

Entrez Gene 22591 MouseEntrez Gene 312560 RatEntrez Gene 7508 Human

Q01831

Background: This gene encodes a component of the nucleotide excision repair (NER) pathway. There are

multiple components involved in the NER pathway, including Xeroderma pigmentosum (XP) A-G and V, Cockayne syndrome (CS) A and B, and trichothiodystrophy (TTD) group A, etc. This component, XPC, plays an important role in the early steps of global genome NER, especially in damage recognition, open complex formation, and repair protein complex formation. Mutations in this gene or some other NER components result in Xeroderma pigmentosum, a rare autosomal recessive disorder characterized by increased sensitivity to sunlight with the development of carcinomas at an early age. Alternatively spliced transcript variants have

been found for this gene. [provided by RefSeq, Mar 2009]



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: p125; RAD4; XP3; XPCC


Protein Families: Druggable Genome

Protein Pathways: Nucleotide excision repair

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY XPC (Cat# [RC204483], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XPC rabbit polyclonal antibody (Cat# [TA890156]).



Western blot analysis of extracts (35ug) from different cell lines and tissues by using anti-XPC rabbit polyclonal antibody.