

Product datasheet for **TB408557**


CD163 CytoSection

Product data:


Product Type:	CytoSections
Description:	Transient overexpression of CD163 (NM_004244), transcript variant 1, in HEK293T cells, paraffin embedded controls for ICC/IHC staining
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC208557
Tag:	C-MYC/DDK
Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Target Detection Antibodies:	CD163 Mouse Monoclonal Antibody [Clone ID: OTI2G12] (TA506380)
ACCN:	NM_004244 , NP_004235
Synonyms:	M130; MM130; SCAR1
Storage:	Room Temperature, or 2-8°C for long term storage
Stability:	Blocks are guaranteed for a year from the date of receipt if proper storage instructions were followed.
Preparation:	HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin.
Note:	This product is for research use only and is not approved for use in humans or in clinical diagnosis.
RefSeq:	NP_004235
Locus ID:	9332
Cytogenetics:	12p13.31
Protein Families:	Druggable Genome, Secreted Protein, Transmembrane




[View online »](#)

Product images:


H&E staining of CD163 overexpressed cytosection (TS408557), HEK293T cells were transfected with cDNA clone [RC208557], 4um sections, 10x magnification



H&E staining of CD163 overexpressed cytosection (TS408557), HEK293T cells were transfected with cDNA clone [RC208557], 4um sections, 40x magnification



Immunohistology staining on CD163 overexpressed cytosection ([TS408557]) with mouse anti DDK clone 11C3 C/N [TA180144] at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen C/N [D22-18]. Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC208557], 4um sections, 40x magnification



Immunohistology staining on HEK293T untransfected control section screened with mouse anti-DDK clone 11C3 C/N [TA180144] at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen C/N [D22-18]. No brown chromogen showing negative stain.