

## **Product datasheet for TF320474**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PKN2 Human shRNA Plasmid Kit (Locus ID 5586)

**Product data:** 

**Product Type:** shRNA Plasmids

**Product Name:** PKN2 Human shRNA Plasmid Kit (Locus ID 5586)

Locus ID: 5586

Synonyms: Pak-2; PAK2; PRKCL2; PRO2042; STK7

**Vector:** pRFP-C-RS (TR30014)

E. coli Selection: Chloramphenicol (34 ug/ml)

**Mammalian Cell** 

Selection:

Puromycin

Format: Retroviral plasmids

**Components:** PKN2 - Human, 4 unique 29mer shRNA constructs in retroviral RFP vector(Gene ID = 5586).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pRFP-C-RS Vector, TR30015, included for free.

**RefSeq:** NM 001320707, NM 001320708, NM 001320709, NM 006256, NM 006256.3, BC006993,

BC029749, BC038832, BC046146, BC062620, BC125199, NM 006256.4

UniProt ID: Q16513

**Summary:** PKC-related serine/threonine-protein kinase and Rho/Rac effector protein that participates in

specific signal transduction responses in the cell. Plays a role in the regulation of cell cycle progression, actin cytoskeleton assembly, cell migration, cell adhesion, tumor cell invasion and transcription activation signaling processes. Phosphorylates CTTN in hyaluronan-induced

astrocytes and hence decreases CTTN ability to associate with filamentous actin.

Phosphorylates HDAC5, therefore lead to impair HDAC5 import. Direct RhoA target required for the regulation of the maturation of primordial junctions into apical junction formation in


bronchial epithelial cells. Required for G2/M phases of the cell cycle progression and abscission during cytokinesis in a ECT2-dependent manner. Stimulates FYN kinase activity

that is required for establishment of skin cell-cell adhesion during keratinocytes

differentiation. Regulates epithelial bladder cells speed and direction of movement during cell migration and tumor cell invasion. Inhibits Akt pro-survival-induced kinase activity. Mediates Rho protein-induced transcriptional activation via the c-fos serum response factor (SRF). Involved in the negative regulation of ciliogenesis (PubMed:27104747).[UniProtKB/Swiss-Prot

Function]







shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.

Performance Guaranteed: OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).