

## **Product datasheet for TF503444**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Syne1 Mouse shRNA Plasmid (Locus ID 64009)

**Product data:** 

**Product Type:** shRNA Plasmids

**Product Name:** Syne1 Mouse shRNA Plasmid (Locus ID 64009)

**Locus ID:** 64009

**Synonyms:** 8B; A330049M09Rik; BE692247; C130039F11Rik; CPG2; mKIAA1756; myne-1; Myne1

**Vector:** pRFP-C-RS (TR30014)

E. coli Selection: Chloramphenicol (34 ug/ml)

Mammalian Cell

Puromycin

Selection:

Format: Retroviral plasmids

**Components:** Syne1 - Mouse, 4 unique 29mer shRNA constructs in retroviral RFP vector(Gene ID = 64009).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pRFP-C-RS Vector, TR30015, included for free.

RefSeq: NM 001079686, NM 001347711, NM 001347732, NM 022027, NM 153399, NM 022027.1,

NM 022027.2, NM 001079686.1, NM 153399.1, NM 153399.2, BC041779, BC054456,

BC110426, BC110428, BC111034

UniProt ID: Q6ZWR6



## Summary:

Multi-isomeric modular protein which forms a linking network between organelles and the actin cytoskeleton to maintain the subcellular spatial organization. As a component of the LINC (Linker of Nucleoskeleton and Cytoskeleton) complex involved in the connection between the nuclear lamina and the cytoskeleton. The nucleocytoplasmic interactions established by the LINC complex play an important role in the transmission of mechanical forces across the nuclear envelope and in nuclear movement and positioning. May be involved in nucleus-centrosome attachment. During interkinetic nuclear migration (INM) at G2 phase and nuclear migration in neural progenitors its LINC complex association with SUN1/2 and probably association with cytoplasmic dynein-dynactin motor complexes functions to pull the nucleus toward the centrosome; SYNE1 and SYNE2 seem to act redundantly in cerebellum, midbrain, brain stem, and other brain regions except cerebral cortex and hippocampus. Required for centrosome migration to the apical cell surface during early ciliogenesis. May be involved in nuclear remodeling during sperm head formation in spermatogenenis; a probable SUN3:SYNE1/KASH1 LINC complex may tether spermatid nuclei to posterior cytoskeletal structures such as the manchette.[UniProtKB/Swiss-Prot Function]

shRNA Design:

Performance Guaranteed:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).