

## Product datasheet for **TG320260**

### AKT1 Human shRNA Plasmid Kit (Locus ID 207)

#### Product data:

Product Type:	shRNA Plasmids
Product Name:	AKT1 Human shRNA Plasmid Kit (Locus ID 207)
Locus ID:	207
Synonyms:	AKT; PKB; PKB-ALPHA; PRKBA; RAC; RAC-ALPHA
Vector:	pGFP-V-RS (TR30007)
E. coli Selection:	Kanamycin
Mammalian Cell Selection:	Puromycin
Format:	Retroviral plasmids
Components:	AKT1 - Human, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 207). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.
RefSeq:	<a href="#">NM_001014431</a> , <a href="#">NM_001014432</a> , <a href="#">NM_005163</a> , <a href="#">NM_001014431.1</a> , <a href="#">NM_005163.1</a> , <a href="#">NM_005163.2</a> , <a href="#">NM_001014432.1</a> , <a href="#">BC000479</a> , <a href="#">BC084538</a> , <a href="#">BC001737</a> , <a href="#">BC063408</a> , <a href="#">NM_001014431.2</a> , <a href="#">NM_001014432.2</a>
UniProt ID:	<a href="#">P31749</a>



[View online »](#)

**Summary:**

This gene encodes one of the three members of the human AKT serine-threonine protein kinase family which are often referred to as protein kinase B alpha, beta, and gamma. These highly similar AKT proteins all have an N-terminal pleckstrin homology domain, a serine/threonine-specific kinase domain and a C-terminal regulatory domain. These proteins are phosphorylated by phosphoinositide 3-kinase (PI3K). AKT/PI3K forms a key component of many signalling pathways that involve the binding of membrane-bound ligands such as receptor tyrosine kinases, G-protein coupled receptors, and integrin-linked kinase. These AKT proteins therefore regulate a wide variety of cellular functions including cell proliferation, survival, metabolism, and angiogenesis in both normal and malignant cells. AKT proteins are recruited to the cell membrane by phosphatidylinositol 3,4,5-trisphosphate (PIP3) after phosphorylation of phosphatidylinositol 4,5-bisphosphate (PIP2) by PI3K. Subsequent phosphorylation of both threonine residue 308 and serine residue 473 is required for full activation of the AKT1 protein encoded by this gene. Phosphorylation of additional residues also occurs, for example, in response to insulin growth factor-1 and epidermal growth factor. Protein phosphatases act as negative regulators of AKT proteins by dephosphorylating AKT or PIP3. The PI3K/AKT signalling pathway is crucial for tumor cell survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating AKT1 which then phosphorylates and inactivates components of the apoptotic machinery. AKT proteins also participate in the mammalian target of rapamycin (mTOR) signalling pathway which controls the assembly of the eukaryotic translation initiation factor 4F (eIF4E) complex and this pathway, in addition to responding to extracellular signals from growth factors and cytokines, is dysregulated in many cancers. Mutations in this gene are associated with multiple types of cancer and excessive tissue growth including Proteus syndrome and Cowden syndrome 6, and breast, colorectal, and ovarian cancers. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2020]

**shRNA Design:**

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact [techsupport@origene.com](mailto:techsupport@origene.com). If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).

**Performance Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).