

# Product datasheet for TG320325

## E2F1 Human shRNA Plasmid Kit (Locus ID 1869)

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

<ul> <li>E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), transactivation domain which is embedded within the transactivation domain. This protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]</li> <li>shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against multiple splice variants at this gene locus. Televistical designed against mu</li></ul>		
Locus ID:1869Synonyms:E2F-1; RBAP1; RBBP3; RBP3Vector:pGFP-V-RS (TR30007)E. coli Selection:KanamycinMammalian CellPuromycinSelection:Retroviral plasmidsComponents:E2F1 - Human, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 1869). Sµg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225.1, NM 005225.2, BC050369, BC050369.2, BC005098, BC058902, NM 005225.3UniProt ID:Q01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor protein and anstick and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. The be certain that your variant of interest is targeted, please contact techsupport@orgene.com	Product Type:	shRNA Plasmids
Synonyms:E2F-1; RBAP1; RBBP3; RBP3Vector:pGFP-V-RS (TR30007)E. coli Selection:KanamycinMammalian CellPuromycinSelection:Retroviral plasmidsFormat:Retroviral plasmidsComponents:E2F1 - Human, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 1869). Sµg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225, NM 005225.1, NM 005225.2, BC050369, BC050369, 2, BC005098, BC058902, NM 005225.3UniProt ID:Q01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor protein a association domain which is embedded within the transcription factor protein a association domain which is erferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. The be certain that your variant of interest is targeted, please contact techsupport@orgene.com	Product Name:	E2F1 Human shRNA Plasmid Kit (Locus ID 1869)
Vector:pGFP-V-RS (TR30007)E. coli Selection:KanamycinMammalian CellPuromycinSelection:Format:Format:Retroviral plasmidsComponents:E2F1 - Human, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 1869). Sµg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225, NM 005225.1, NM 005225.2, BC050369, BC050369, 2, BC050398, BC058902, NM 005225.3UniProt ID:Q01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor protein association domain which is embedded within the transactivation domain. This protein an association domain which is embedded within the transactivation domain. This protein an association domain which is embedded within the transactivation domain. This protein an association domain which is embedded within the transactivation domain. This protein an association domain which is embedded within the transactivation domain. This protein an another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein an another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein an another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein an another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein an another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein an another 2 members, E2F2 and	Locus ID:	1869
E. coli Selection:KanamycinMammalian Cell Selection:PuromycinFormat:Retroviral plasmidsComponents:E2F1 - Human, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 1869). Syg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225, NM 005225.1, NM 005225.2, BC050369, BC050369.2, BC005098, BC058902, NM 005225.3UniProt ID:O01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor protein and association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein and mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq. Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. The be certain that your variant of interest is targeted, please contact techsupport@origene.com	Synonyms:	E2F-1; RBAP1; RBBP3; RBP3
Mammalian Cell Selection:PuromycinFormat:Retroviral plasmidsComponents:E2F1 - Human, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 1869). Sµg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225, NM 005225,1, NM 005225,2, BC050369, BC050369,2, BC05098, BC058902, NM 005225,3UniProt ID:Q01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins and association domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein and mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. The be certain that your variant of interest is targeted, please contact techsupport@origene.com	Vector:	pGFP-V-RS (TR30007)
Selection:Format:Retroviral plasmidsComponents:E2F1 - Human, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 1869). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225, NM 005225.1, NM 005225.2, BC050369, BC050369.2, BC005098, BC058902, NM 005225.3UniProt ID:Q01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor protein and ansociation domain which is embedded within the transactivation domain. This protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. The be certain that your variant of interest is targeted, please contact techsupport@origene.com	E. coli Selection:	Kanamycin
Components:E2F1 - Hunan, 4 unique 29mer shRNA constructs in retroviral GFP vector(Gene ID = 1869). Sµg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225, NM 005225.1, NM 005225.2, BC050369, BC050369.2, BC005098, BC058902, NM 005225.3UniProt ID:O01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), transactivation domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. The be certain that your variant of interest is targeted, please contact techsupport@origene.com		Puromycin
Sug purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-V-RS Vector, TR30013, included for free.RefSeq:NM 005225, NM 005225.1, NM 005225.2, BC050369, BC050369.2, BC005098, BC058902, NM 005225.3UniProt ID:Q01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com	Format:	Retroviral plasmids
NM 005225.3UniProt ID:O01094Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com	Components:	5µg purified plasmid DNA per construct
Summary:The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com	RefSeq:	
<ul> <li>E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of th family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]</li> <li>shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. The be certain that your variant of interest is targeted, please contact techsupport@origene.com</li> </ul>	UniProt ID:	<u>Q01094</u>
be certain that your variant of interest is targeted, please contact techsupport@origene.con	Summary:	proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by
	shRNA Design:	These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u> . If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u> .



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **CRIGENE** E2F1 Human shRNA Plasmid Kit (Locus ID 1869) – TG320325

Performance Guaranteed: OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US