

## Product datasheet for **TL302335**

### PDP1 Human shRNA Plasmid Kit (Locus ID 54704)

#### Product data:

Product Type:	shRNA Plasmids
Product Name:	PDP1 Human shRNA Plasmid Kit (Locus ID 54704)
Locus ID:	54704
Synonyms:	PDH; PDP; PDPC; PPM2C
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	PDP1 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 54704). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	<a href="#">NM_001161778</a> , <a href="#">NM_001161779</a> , <a href="#">NM_001161780</a> , <a href="#">NM_001161781</a> , <a href="#">NM_018444</a> , <a href="#">NM_018444.1</a> , <a href="#">NM_018444.2</a> , <a href="#">NM_018444.3</a> , <a href="#">NM_001161779.1</a> , <a href="#">NM_001161780.1</a> , <a href="#">NM_001161778.1</a> , <a href="#">NM_001161781.1</a> , <a href="#">BC047619</a> , <a href="#">BC047619.1</a> , <a href="#">BC064978</a> , <a href="#">BC098343</a> , <a href="#">BM973633</a> , <a href="#">NM_001161781.2</a> , <a href="#">NM_018444.4</a> , <a href="#">NM_001161780.2</a>
UniProt ID:	<a href="#">Q9P0J1</a>
Summary:	Pyruvate dehydrogenase (E1) is one of the three components (E1, E2, and E3) of the large pyruvate dehydrogenase complex. Pyruvate dehydrogenase kinases catalyze phosphorylation of serine residues of E1 to inactivate the E1 component and inhibit the complex. Pyruvate dehydrogenase phosphatases catalyze the dephosphorylation and activation of the E1 component to reverse the effects of pyruvate dehydrogenase kinases. Pyruvate dehydrogenase phosphatase is a heterodimer consisting of catalytic and regulatory subunits. Two catalytic subunits have been reported; one is predominantly expressed in skeletal muscle and another one is much more abundant in the liver. The catalytic subunit, encoded by this gene, is the former, and belongs to the protein phosphatase 2C (PP2C) superfamily. Along with the pyruvate dehydrogenase complex and pyruvate dehydrogenase kinases, this enzyme is located in the mitochondrial matrix. Mutation in this gene causes pyruvate dehydrogenase phosphatase deficiency. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.[provided by RefSeq, Jun 2009]



[View online »](#)

- shRNA Design:** These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact [techsupport@origene.com](mailto:techsupport@origene.com). If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).
- Performance Guaranteed:** OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.
- For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).