

## Product datasheet for **TL302749**

### OSBPL9 Human shRNA Plasmid Kit (Locus ID 114883)

#### Product data:

Product Type:	shRNA Plasmids
Product Name:	OSBPL9 Human shRNA Plasmid Kit (Locus ID 114883)
Locus ID:	114883
Synonyms:	ORP-9; ORP9
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	OSBPL9 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 114883). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	<a href="#">NM_001330580</a> , <a href="#">NM_024586</a> , <a href="#">NM_148904</a> , <a href="#">NM_148905</a> , <a href="#">NM_148906</a> , <a href="#">NM_148907</a> , <a href="#">NM_148908</a> , <a href="#">NM_148909</a> , <a href="#">NR_036662</a> , <a href="#">NM_001350208</a> , <a href="#">NM_001350209</a> , <a href="#">NM_001350210</a> , <a href="#">NM_148904.1</a> , <a href="#">NM_148904.2</a> , <a href="#">NM_148904.3</a> , <a href="#">NM_148909.1</a> , <a href="#">NM_148909.2</a> , <a href="#">NM_148909.3</a> , <a href="#">NM_148905.1</a> , <a href="#">NM_148905.2</a> , <a href="#">NM_148905.3</a> , <a href="#">NM_148906.1</a> , <a href="#">NM_148906.2</a> , <a href="#">NM_148908.1</a> , <a href="#">NM_148908.2</a> , <a href="#">NM_148908.3</a> , <a href="#">NM_148907.1</a> , <a href="#">NM_148907.2</a> , <a href="#">NM_024586.1</a> , <a href="#">NM_024586.2</a> , <a href="#">NM_024586.3</a> , <a href="#">NM_024586.4</a> , <a href="#">NM_024586.5</a> , <a href="#">BC025978</a> , <a href="#">BC025978.2</a> , <a href="#">BC006160</a> , <a href="#">NM_148904.4</a> , <a href="#">NM_148908.4</a> , <a href="#">NM_148907.3</a> , <a href="#">NM_148905.4</a> , <a href="#">NM_148909.4</a> , <a href="#">NM_024586.6</a>
UniProt ID:	<a href="#">Q96SU4</a>
Summary:	This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors. Most members contain an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain, although some members contain only the sterol-binding domain. This family member functions as a cholesterol transfer protein that regulates Golgi structure and function. Multiple transcript variants, most of which encode distinct isoforms, have been identified. Related pseudogenes have been identified on chromosomes 3, 11 and 12. [provided by RefSeq, Jul 2010]
shRNA Design:	These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a> . If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a> .



[View online »](#)

**Performance  
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).