

## Product datasheet for **TL302968**

### **NFX1 Human shRNA Plasmid Kit (Locus ID 4799)**

#### **Product data:**

|                                  |   |
|----------------------------------|---|
| <b>Product Type:</b>             | shRNA Plasmids  |
| <b>Product Name:</b>             | NFX1 Human shRNA Plasmid Kit (Locus ID 4799)  |
| <b>Locus ID:</b>                 | 4799  |
| <b>Synonyms:</b>                 | NFX2; TEG-42; Tex42   |
| <b>Vector:</b>                   | pGFP-C-shLenti (TR30023)  |
| <b>E. coli Selection:</b>        | Chloramphenicol (34 ug/ml)  |
| <b>Mammalian Cell Selection:</b> | Puromycin   |
| <b>Format:</b>                   | Lentiviral plasmids   |
| <b>Components:</b>               | NFX1 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 4799).<br>5µg purified plasmid DNA per construct<br>29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.  |
| <b>RefSeq:</b>                   | <a href="#">NM_001318758</a> , <a href="#">NM_002504</a> , <a href="#">NM_147133</a> , <a href="#">NM_147134</a> , <a href="#">NM_002504.1</a> , <a href="#">NM_002504.2</a> ,<br><a href="#">NM_002504.3</a> , <a href="#">NM_002504.4</a> , <a href="#">NM_002504.5</a> , <a href="#">NM_147133.1</a> , <a href="#">NM_147133.2</a> , <a href="#">NM_147134.2</a> ,<br><a href="#">NM_147134.3</a> , <a href="#">BC012151</a> , <a href="#">BC012151.1</a> , <a href="#">BC006447</a> , <a href="#">NM_147134.4</a> , <a href="#">NM_002504.6</a>   |
| <b>UniProt ID:</b>               | <a href="#">Q12986</a>  |
| <b>Summary:</b>                  | MHC class II gene expression is controlled primarily at the transcriptional level by transcription factors that bind to the X and Y boxes, two highly conserved elements in the proximal promoter of MHC class II genes. The protein encoded by this gene is a transcriptional repressor capable of binding to the conserved X box motif of HLA-DRA and other MHC class II genes in vitro. The protein may play a role in regulating the duration of an inflammatory response by limiting the period in which class II MHC molecules are induced by IFN-gamma. Three alternative splice variants, each of which encodes a different isoform, have been identified. [provided by RefSeq, Jul 2008] |
| <b>shRNA Design:</b>             | These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a> . If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a> .  |



[View online »](#)

**Performance  
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).