

## **Product datasheet for TL303587**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## LAMC2 Human shRNA Plasmid Kit (Locus ID 3918)

**Product data:** 

**Product Type:** shRNA Plasmids

Product Name: LAMC2 Human shRNA Plasmid Kit (Locus ID 3918)

**Locus ID:** 3918

Synonyms: B2T; BM600; CSF; EBR2; EBR2A; LAMB2T; LAMNB2

**Vector:** pGFP-C-shLenti (TR30023)

E. coli Selection: Chloramphenicol (34 ug/ml)

Mammalian Cell

Puromycin

Selection:

Format: Lentiviral plasmids

**Components:** LAMC2 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 3918).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.

RefSeq: NM 005562, NM 018891, NM 018891.1, NM 018891.2, NM 005562.1, NM 005562.2,

BC112286, BC113378, BC143653, NM 018891.3

UniProt ID: Q13753



Summary:

Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminins, composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively), have a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. Several isoforms of each chain have been described. Different alpha, beta and gamma chain isomers combine to give rise to different heterotrimeric laminin isoforms which are designated by Arabic numerals in the order of their discovery, i.e. alpha1beta1gamma1 heterotrimer is laminin 1. The biological functions of the different chains and trimer molecules are largely unknown, but some of the chains have been shown to differ with respect to their tissue distribution, presumably reflecting diverse functions in vivo. This gene encodes the gamma chain isoform laminin, gamma 2. The gamma 2 chain, formerly thought to be a truncated version of beta chain (B2t), is highly homologous to the gamma 1 chain; however, it lacks domain VI, and domains V, IV and III are shorter. It is expressed in several fetal tissues but differently from gamma 1, and is specifically localized to epithelial cells in skin, lung and kidney. The gamma 2 chain together with alpha 3 and beta 3 chains constitute laminin 5 (earlier known as kalinin), which is an integral part of the anchoring filaments that connect epithelial cells to the underlying basement membrane. The epithelium-specific expression of the gamma 2 chain implied its role as an epithelium attachment molecule, and mutations in this gene have been associated with junctional epidermolysis bullosa, a skin disease characterized by blisters due to disruption of the epidermal-dermal junction. Two transcript variants resulting from alternative splicing of the 3' terminal exon, and encoding different isoforms of gamma 2 chain, have been described. The two variants are differentially expressed in embryonic tissues, however, the biological significance of the two forms is not known. Transcript variants utilizing alternative polyA\_signal have also been noted in literature. [provided by RefSeq, Aug 2011]

shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.



## Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).