

## Product datasheet for **TL305184**

### **BORIS (CTCFL) Human shRNA Plasmid Kit (Locus ID 140690)**

#### **Product data:**

Product Type:	shRNA Plasmids
Product Name:	BORIS (CTCFL) Human shRNA Plasmid Kit (Locus ID 140690)
Locus ID:	140690
Synonyms:	BORIS; CT27; CTCF-T; dj579F20.2; HMGB1L1
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	CTCFL - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 140690). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	<a href="#">NM_001269040</a> , <a href="#">NM_001269041</a> , <a href="#">NM_001269042</a> , <a href="#">NM_001269043</a> , <a href="#">NM_001269044</a> , <a href="#">NM_001269045</a> , <a href="#">NM_001269046</a> , <a href="#">NM_001269047</a> , <a href="#">NM_001269048</a> , <a href="#">NM_001269049</a> , <a href="#">NM_001269050</a> , <a href="#">NM_001269051</a> , <a href="#">NM_001269052</a> , <a href="#">NM_001269054</a> , <a href="#">NM_001269055</a> , <a href="#">NM_080618</a> , <a href="#">NR_072975</a> , <a href="#">NM_080618.1</a> , <a href="#">NM_080618.2</a> , <a href="#">NM_080618.3</a> , <a href="#">NM_001269055.1</a> , <a href="#">NM_001269051.1</a> , <a href="#">NM_001269052.1</a> , <a href="#">NM_001269050.1</a> , <a href="#">NM_001269047.1</a> , <a href="#">NM_001269054.1</a> , <a href="#">NM_001269049.1</a> , <a href="#">NM_001269048.1</a> , <a href="#">NM_001269046.1</a> , <a href="#">NM_001269045.1</a> , <a href="#">NM_001269042.1</a> , <a href="#">NM_001269040.1</a> , <a href="#">NM_001269041.1</a> , <a href="#">NM_001269044.1</a> , <a href="#">NM_001269043.1</a> , <a href="#">BC060763</a> , <a href="#">BC130486</a> , <a href="#">BC137482</a> , <a href="#">BC137483</a> , <a href="#">NM_001269055.2</a> , <a href="#">NM_001269050.2</a>
UniProt ID:	<a href="#">Q8NI51</a>
Summary:	CCCTC-binding factor (CTCF), an 11-zinc-finger factor involved in gene regulation, utilizes different zinc fingers to bind varying DNA target sites. CTCF forms methylation-sensitive insulators that regulate X-chromosome inactivation. This gene is a paralog of CTCF and appears to be expressed primarily in the cytoplasm of spermatocytes, unlike CTCF which is expressed primarily in the nucleus of somatic cells. CTCF and the protein encoded by this gene are normally expressed in a mutually exclusive pattern that correlates with resetting of methylation marks during male germ cell differentiation. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2012]



[View online »](#)

**shRNA Design:**

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact [techsupport@origene.com](mailto:techsupport@origene.com). If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).

**Performance Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).