

Product datasheet for **TL306422**

Bim (BCL2L11) Human shRNA Plasmid Kit (Locus ID 10018)


Product data:

Product Type:	shRNA Plasmids
Product Name:	Bim (BCL2L11) Human shRNA Plasmid Kit (Locus ID 10018)
Locus ID:	10018
Synonyms:	BAM; BIM; BOD
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	BCL2L11 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 10018). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	NM_001204106 , NM_001204107 , NM_001204108 , NM_001204109 , NM_001204110 , NM_001204111 , NM_001204112 , NM_001204113 , NM_006538 , NM_138621 , NM_138622 , NM_138623 , NM_138624 , NM_138625 , NM_138626 , NM_138627 , NM_207002 , NM_207003 , NM_138621.1 , NM_138621.2 , NM_138621.3 , NM_138621.4 , NM_006538.2 , NM_006538.3 , NM_006538.4 , NM_207002.1 , NM_207002.2 , NM_207002.3 , NM_138627.1 , NM_138627.2 , NM_138627.3 , NM_001204111.1 , NM_138625.1 , NM_138625.2 , NM_138625.3 , NM_207003.1 , NM_207003.2 , NM_001204112.1 , NM_001204110.1 , NM_001204107.1 , NM_001204113.1 , NM_001204106.1 , NM_138623.1 , NM_138623.2 , NM_138623.3 , NM_138624.1 , NM_138624.3 , NM_138626.1 , NM_138626.2 , NM_138626.3 , NM_001204109.1 , NM_138622.1 , NM_138622.2 , NM_138622.3 , NM_001204108.1 , BC033694 , BC033694.1 , NM_006538.5 , NM_001204106.2 , NM_001204110.2 , NM_138625.4 , NM_138627.4 , NM_138621.5 , NM_207003.3 , NM_001204111.2 , NM_001204112.2
UniProt ID:	O43521




[View online »](#)


- Summary:** The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The protein encoded by this gene contains a Bcl-2 homology domain 3 (BH3). It has been shown to interact with other members of the BCL-2 protein family and to act as an apoptotic activator. The expression of this gene can be induced by nerve growth factor (NGF), as well as by the forkhead transcription factor FKHL1, which suggests a role of this gene in neuronal and lymphocyte apoptosis. Transgenic studies of the mouse counterpart suggested that this gene functions as an essential initiator of apoptosis in thymocyte-negative selection. Several alternatively spliced transcript variants of this gene have been identified. [provided by RefSeq, Jun 2013]
- shRNA Design:** These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).
- Performance Guaranteed:** OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.
- For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:


GFP signal was observed under microscope at 48 hours after transduction of TL306422A virus into HEK293 cells. TL306422A virus was prepared using lenti-shRNA TL306422A and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of TL306422B virus into HEK293 cells. TL306422B virus was prepared using lenti-shRNA TL306422B and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL306422C] virus into HEK293 cells. [TL306422C] virus was prepared using lenti-shRNA [TL306422C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL306422D] virus into HEK293 cells. [TL306422D] virus was prepared using lenti-shRNA [TL306422D] and [TR30037] packaging kit.