

Product datasheet for **TL306881**

ACE2 Human shRNA Plasmid Kit (Locus ID 59272) (Angiotensin Converting Enzyme 2)

Product data:

| | |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Type: | shRNA Plasmids |
| Product Name: | ACE2 Human shRNA Plasmid Kit (Locus ID 59272) (Angiotensin Converting Enzyme 2) |
| Locus ID: | 59272 |
| Synonyms: | ACEH |
| Vector: | pGFP-C-shLenti (TR30023) |
| E. coli Selection: | Chloramphenicol (34 ug/ml) |
| Mammalian Cell Selection: | Puromycin |
| Format: | Lentiviral plasmids |
| Components: | ACE2 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 59272). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free. |
| RefSeq: | NM_021804 , NM_021804.1 , NM_021804.2 , BC048094 , BC048094.1 , BC032938 , BC039902 , BC048144 , BC059378 |
| UniProt ID: | Q9BYF1 |
| Summary: | The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. ACE2 is known to be expressed in various human organs, and its organ- and cell-specific expression suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronavirus HCoV-NL63 and the human severe acute respiratory syndrome coronaviruses, SARS-CoV and SARS-CoV-2, the latter is the causative agent of coronavirus disease-2019 (COVID-19). Multiple splice variants have been found for this gene and the dACE2 (or MIRb-ACE2) splice variant has been found to be interferon inducible. [provided by RefSeq, Nov 2020] |
| shRNA Design: | These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com . If you need a special design or shRNA sequence, please utilize our custom shRNA service . |




[View online »](#)


**Performance
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:


GFP signal was observed under microscope at 48 hours after transduction of TL306881A virus into HEK293 cells. TL306881A virus was prepared using lenti-shRNA TL306881A and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of TL306881B virus into HEK293 cells. TL306881B virus was prepared using lenti-shRNA TL306881B and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL306881C] virus into HEK293 cells. [TL306881C] virus was prepared using lenti-shRNA [TL306881C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL306881D] virus into HEK293 cells. [TL306881D] virus was prepared using lenti-shRNA [TL306881D] and [TR30037] packaging kit.