

Product datasheet for **TL309351V**

FATP2 (SLC27A2) Human shRNA Lentiviral Particle (Locus ID 11001)

Product data:

| | |
|---------------|--|
| Product Type: | shRNA Lentiviral Particles |
| Product Name: | FATP2 (SLC27A2) Human shRNA Lentiviral Particle (Locus ID 11001) |
| Locus ID: | 11001 |
| Synonyms: | ACSVL1; FACVL1; FATP2; hFACVL1; HsT17226; VLACS; VLCS |
| Vector: | pGFP-C-shLenti (TR30023) |
| Format: | Lentiviral particles |
| Components: | SLC27A2 - Human shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1 scramble control), 0.5 ml each, >10 ⁷ TU/ml. |
| RefSeq: | NM_001159629 , NM_003645 , NM_003645.2 , NM_003645.3 , NM_001159629.1 , BC057770 , BM980042 , NM_003645.4 |
| UniProt ID: | O14975 |
| Summary: | The protein encoded by this gene is an isozyme of long-chain fatty-acid-coenzyme A ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme activates long-chain, branched-chain and very-long-chain fatty acids containing 22 or more carbons to their CoA derivatives. It is expressed primarily in liver and kidney, and is present in both endoplasmic reticulum and peroxisomes, but not in mitochondria. Its decreased peroxisomal enzyme activity is in part responsible for the biochemical pathology in X-linked adrenoleukodystrophy. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2009] |
| shRNA Design: | These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com . If you need a special design or shRNA sequence, please utilize our custom shRNA service . |




[View online »](#)


**Performance
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.


For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

Product images:


GFP signal was observed under microscope at 48 hours after transduction of TL309351A virus into HEK293 cells. TL309351A virus was prepared using lenti-shRNA TL309351A and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of TL309351B virus into HEK293 cells. TL309351B virus was prepared using lenti-shRNA TL309351B and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL309351C] virus into HEK293 cells. [TL309351C] virus was prepared using lenti-shRNA [TL309351C] and [TR30037] packaging kit.



GFP signal was observed under microscope at 48 hours after transduction of [TL309351D] virus into HEK293 cells. [TL309351D] virus was prepared using lenti-shRNA [TL309351D] and [TR30037] packaging kit.