

Product datasheet for **TL312000**

KCNMA1 Human shRNA Plasmid Kit (Locus ID 3778)

Product data:

Product Type:	shRNA Plasmids
Product Name:	KCNMA1 Human shRNA Plasmid Kit (Locus ID 3778)
Locus ID:	3778
Synonyms:	bA205K10.1; BKTM; CADEDS; hSlo; IEG16; KCa1.1; LIWAS; MaxiK; mSLO1; PNKD3; SAKCA; SLO; SLO-ALPHA; SLO1
Vector:	pGFP-C-shLenti (TR30023)
E. coli Selection:	Chloramphenicol (34 ug/ml)
Mammalian Cell Selection:	Puromycin
Format:	Lentiviral plasmids
Components:	KCNMA1 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 3778). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.
RefSeq:	<u>BC024965</u> , <u>NM_001014797</u> , <u>NM_001161352</u> , <u>NM_001161353</u> , <u>NM_001271518</u> , <u>NM_001271519</u> , <u>NM_001271520</u> , <u>NM_001271521</u> , <u>NM_001271522</u> , <u>NM_001322829</u> , <u>NM_001322830</u> , <u>NM_001322832</u> , <u>NM_001322835</u> , <u>NM_001322836</u> , <u>NM_001322837</u> , <u>NM_001322838</u> , <u>NM_001322839</u> , <u>NM_002247</u> , <u>NM_002247.1</u> , <u>NM_002247.2</u> , <u>NM_002247.3</u> , <u>NM_001014797.1</u> , <u>NM_001014797.2</u> , <u>NM_001161353.1</u> , <u>NM_001161352.1</u> , <u>NM_001271522.1</u> , <u>NM_001271520.1</u> , <u>NM_001271518.1</u> , <u>NM_001271519.1</u> , <u>NM_001271521.1</u> , <u>BC009695</u> , <u>BC062659</u> , <u>BC137115</u> , <u>BC137137</u> , <u>BC144496</u> , <u>NM_001271518.2</u> , <u>NM_001271519.2</u> , <u>NM_001271520.2</u> , <u>NM_001161352.2</u> , <u>NM_001161353.2</u> , <u>NM_001014797.3</u> , <u>NM_001271522.2</u> , <u>NM_002247.4</u> , <u>NM_001271521.2</u>
UniProt ID:	<u>Q12791</u>
Summary:	MaxiK channels are large conductance, voltage and calcium-sensitive potassium channels which are fundamental to the control of smooth muscle tone and neuronal excitability. MaxiK channels can be formed by 2 subunits: the pore-forming alpha subunit, which is the product of this gene, and the modulatory beta subunit. Intracellular calcium regulates the physical association between the alpha and beta subunits. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]



[View online »](#)

- shRNA Design:** These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).
- Performance Guaranteed:** OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.
- For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).