

## **Product datasheet for TL313086V**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

#### **FAM20B Human shRNA Lentiviral Particle (Locus ID 9917)**

#### **Product data:**

**Product Type:** shRNA Lentiviral Particles

**Product Name:** FAM20B Human shRNA Lentiviral Particle (Locus ID 9917)

Locus ID: 9917 Synonyms: gxk1

**Vector:** pGFP-C-shLenti (TR30023)

Format: Lentiviral particles

Components: FAM20B - Human shRNA lentiviral particles (4 unique 29mer target-specific shRNA, 1

scramble control), 0.5 ml each, >10^7 TU/ml.

RefSeq: NM 014864, NM 001324310, NM 001324311, NM 014864.1, NM 014864.2, NM 014864.3,

BC051794, BC051794.1, BC046441

UniProt ID: <u>075063</u>

**Summary:** Responsible for the 2-O-phosphorylation of xylose in the glycosaminoglycan-protein linkage

region of proteoglycans thereby regulating the amount of mature GAG chains. Sulfated glycosaminoglycans (GAGs), including heparan sulfate and chondroitin sulfate, are synthesized on the so-called common GAG-protein linkage region (GlcUAbeta1-3Galbeta1-3Galbeta1-4Xylbeta1-O-Ser) of core proteins, which is formed by the stepwise addition of

monosaccharide residues by the respective specific glycosyltransferases. Xylose 2-O-phosphorylation may influence the catalytic activity of B3GAT3 (GlcAT-I) which completes the precursor tetrasaccharide of GAG-protein linkage regions on which the repeating disaccharide

region is synthesized.[UniProtKB/Swiss-Prot Function]

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.







# Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).