

## **Product datasheet for TL313711**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## **CRYBB2 Human shRNA Plasmid Kit (Locus ID 1415)**

**Product data:** 

**Product Type:** shRNA Plasmids

**Product Name:** CRYBB2 Human shRNA Plasmid Kit (Locus ID 1415)

**Locus ID:** 1415

Synonyms: CCA2; CRYB2; CRYB2A; CTRCT3; D22S665

**Vector:** pGFP-C-shLenti (TR30023)

E. coli Selection: Chloramphenicol (34 ug/ml)

Mammalian Cell

Selection:

Puromycin

Format: Lentiviral plasmids

Components: CRYBB2 - Human, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 1415).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.

RefSeq: NM 000496, NM 000496.1, NM 000496.2, BC069535, BC069535.1, BM667912, NM 000496.3

UniProt ID: P43320

**Summary:** Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The

latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Beta-crystallins, the most heterogeneous, differ by the presence of the C-terminal extension (present in the basic group, none in the acidic group). Beta-crystallins form aggregates of different sizes and are able to self-associate to form dimers or to form heterodimers with other beta-crystallins. This gene, a beta basic group member, is part of a gene cluster with beta-A4, beta-B1, and beta-B3. A chain-terminating mutation was found to cause type 2

cerulean cataracts. [provided by RefSeq, Jul 2008]





## CRYBB2 Human shRNA Plasmid Kit (Locus ID 1415) - TL313711

shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.

Performance Guaranteed: OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).